

Low doses of glyphosate enhance growth, CO₂ assimilation, stomatal conductance and transpiration in sugarcane and eucalyptus

Renan F Nascentes,^a Caio A Carbonari,^{a*} Plinio S Simões,^a Marcela C Brunelli,^a Edivaldo D Velini^a and Stephen O Duke^b

Abstract

INTRODUCTION: Sublethal doses of herbicides can enhance plant growth and stimulate other process, an effect known as hormesis. The magnitude of hormesis is dependent on the plant species, the herbicide and its dose, plant development stage and environmental parameters. Glyphosate hormesis is well established, but relatively little is known of the mechanism of this phenomenon. The objective of this study was to determine if low doses of glyphosate that cause growth stimulation in sugarcane and eucalyptus concomitantly stimulate CO₂ assimilation.

RESULTS: Shoot dry weight in both species increased at both 40 and 60 days after application of 6.2 to 20.2 g a.e. ha⁻¹ glyphosate. The level of enhanced shoot dry weight was 11 to 37%, depending on the time after treatment and the species. Concomitantly, CO₂ assimilation, stomatal conductance and transpiration were increased by glyphosate doses similar to those that caused growth increases.

CONCLUSION: Glyphosate applied at low doses increased the dry weight of sugarcane and eucalyptus plants in all experiments. This hormetic effect was related to low dose effects on CO₂ assimilation rate, stomatal conductance and transpiration rate, indicating that low glyphosate doses enhance photosynthesis of plants.

© 2017 Society of Chemical Industry

Keywords: hormesis; *Eucalyptus* sp.; *Saccharum* spp.; low doses

1 INTRODUCTION

Glyphosate is a systemic herbicide that inhibits the shikimic acid pathway enzyme 5-enolpyruvylshikimate 3-phosphate synthase.^{1,2} The shikimic acid pathway is responsible for the production of aromatic amino acids (tryptophan, phenylalanine and tyrosine) and many other phenolic compounds derived from these amino acids.^{3–7}

Hormesis, the stimulatory effect of a subtoxic dose of a toxin,⁸ is commonly found with herbicides.⁹ In particular, there are numerous reports that low, subtoxic doses of glyphosate enhance plant growth and various physiological parameters.⁹ In general, glyphosate hormesis is more reproducible and more pronounced than hormesis associated with other herbicides. For example, glyphosate applications between 1 and 10 g a.e. ha⁻¹ can stimulate growth of eucalyptus (*Eucalyptis grandis*) from 40 to 100% depending on the plant part measured,¹⁰ whereas hormetic effects of this magnitude are rare with other herbicides.⁹

Although utilizing glyphosate hormesis to improve yield is a daunting goal because of the unpredictability of the effect in the field, a better understanding of the mechanism of glyphosate-caused hormesis might improve the predictability of this phenomenon. The mechanism of glyphosate- or any other herbicide-caused hormesis is not clearly understood, although, in the case of glyphosate-caused hormesis, it might be associated partly with the herbicide site of action,¹¹ since this effect is not observed in glyphosate-resistant plants at the doses that cause

hormesis in glyphosate-sensitive plants.¹⁰ Low hormetic doses of glyphosate stimulated photosynthesis in barley plants.¹¹ The same authors proposed that at phytotoxic glyphosate doses, inhibition of photosynthesis is a secondary effect of inhibition of the shikimate pathway.

The objective of the present study was to determine the relationship between glyphosate hormesis, carbon assimilation and transpiration in eucalyptus and sugarcane in an effort to further understand the mechanism of glyphosate hormesis in plants. We chose sugarcane and eucalyptus for these experiments because of our previous success in obtaining robust glyphosate-caused hormesis with these species.^{10,12}

2 MATERIALS AND METHODS

Sugarcane stalks and eucalyptus seedlings were acquired from a commercial plant nursery. Two experiments were conducted in a greenhouse (air temperature of 27.0 ± 3.4 °C, with 65 ± 5% relative

* Correspondence to: CA Carbonari, São Paulo State University (UNESP), Faculty of Agronomic Sciences, Laboratory of Weed Science, Botucatu, SP, CEP 18603-970 PB 237, Brazil. E-mail: carbonari@fca.unesp.br

a São Paulo State University (UNESP), Faculty of Agronomic Sciences, Laboratory of Weed Science, Botucatu, Brazil

b USDA-ARS Natural Products Utilization Research Unit, University, MS, USA




Figure 1. Leaf, stem and total dry weight of sugarcane plants 40 days (A) and 60 days (B) after application of glyphosate. Data and statistics for this figure are provided in Table 1.




Figure 2. Leaf, stem and total dry weight of eucalyptus plants 40 days (A) and 60 days (B) after application of glyphosate. Data and statistics for this figure are provided in Table 2.

humidity, without supplementary lighting) with sugarcane (*Saccharum officinarum*, variety SP801842) and eucalyptus (*Eucalyptus urograndis*, clone 144). In the first experiment the plants were grown until 40 days after glyphosate application (DAA) and in the other until 60 DAA. In the first experiment, chlorophyll content was determined at 15 and 30 DAA, CO_2 assimilation rate, stomatal conductance and transpiration rate were determined at 15 DAA, and dry matter was determined at the end of the experiment (40 DAA). In the second experiment, dry matter and glyphosate and shikimate content were determined 60 DAA. Both species were

cultivated in 5 L pots filled with soil. Soil fertilization was made as indicated by the soil chemical analysis, following technical recommendations for these crops. Water was provided as required for normal growth and development of the plants.

2.1 Glyphosate treatment and growth measurements

Treatments consisted of 0, 1.8, 3.6, 7.2, 18, 36, 72, 180, 360, 720 g a.e. ha^{-1} of glyphosate distributed in a complete randomized experimental design with five replications. A commercial glyphosate isopropylamine 480 g L^{-1} (360 g a.e. L^{-1}) SL (Roundup

Table 1. Parameters for the regression equations and *F* values for leaf, stem and total dry weight of sugarcane plants (40 and 60 DAA). Values represent regression parameters $\pm$ standard error. Curves from these data are provided in Fig. 1

	Dry weight (% of control)		
	40 days after treatment		
	Leaf	Stem	Total
Hypothesis $f \neq 0$	17.51**	13.31**	24.03**
Regression	44.90**	26.60**	48.31**
Model	Model 1	Model 1	Model 1
R²	0.99	0.98	0.99
Constants			
<i>g</i>	-2.33 ± 0.47	-2.42 ± 0.70	-2.37 ± 0.56
<i>b</i>	1.79 ± 0.25	1.62 ± 0.26	1.68 ± 0.23
<i>k</i>	48.37 ± 8.91	54.13 ± 15.33	51.49 ± 11.51
<i>d</i>	50.15 ± 8.83	37.47 ± 13.56	43.22 ± 10.91
<i>f</i>	9.58 ± 6.09	11.70 ± 9.97	10.85 ± 7.73
	60 days after treatment		
	Leaf	Stem	Total
	Leaf	Stem	Total
Hypothesis $f \neq 0$	11.50**	4.33*	4.32*
Regression	102.20**	50.51**	68.42**
Model	Model 1	Model 1	Model 1
R²	0.99	0.99	0.99
Constants			
<i>g</i>	-3.50 ± 0.10	-3.75 ± 0.29	-3.53 ± 0.31
<i>b</i>	3.06 ± 0.33	2.85 ± 0.75	2.62 ± 0.62
<i>k</i>	87.40 ± 3.19	94.09 ± 7.36	91.75 ± 7.88
<i>d</i>	17.71 ± 2.66	19.77 ± 5.12	18.58 ± 3.43
<i>f</i>	2.47 ± 0.66	1.44 ± 1.34	2.13 ± 1.84

* $P < 0.05$; ** $P < 0.01$.

Table 2. Parameters for the regression equations and *F* values for leaf, stem and total dry weight of eucalyptus plants (40 and 60 DAA). Values represent regression parameters $\pm$ standard error. Curves from these data are provided in Fig. 2

	Dry weight (% of control)		
	40 days after treatment		
	Leaf	Stem	Total
Hypothesis $f \neq 0$	8.63**	8.74**	4.55**
Regression	55.8**	7.53**	15.52**
Model	Model 1	Model 1	Model 1
R²	0.99	0.99	0.99
Constants			
<i>g</i>	-1.59 ± 0.73	1.59 ± 0.80	-1.97 ± 0.77
<i>b</i>	1.52 ± 0.24	1.56 ± 0.27	1.48 ± 0.20
<i>k</i>	14.88 ± 11.10	12.10 ± 10.47	32.60 ± 12.24
<i>d</i>	84.96 ± 10.88	86.05 ± 13.42	68.41 ± 13.68
<i>f</i>	18.93 ± 13.60	15.27 ± 12.35	14.16 ± 0.20
	60 days after treatment		
	Leaf	Stem	Total
	Leaf	Stem	Total
Hypothesis $f \neq 0$	5.85*	15.57**	4.75*
Regression	25.47**	5.54**	21.25**
Model	Model 1	Model 1	Model 1
R²	0.99	0.99	0.99
Constants			
<i>g</i>	-3.80 ± 0.58	-1.81 ± 0.47	-3.77 ± 0.74
<i>b</i>	1.95 ± 0.37	1.77 ± 0.30	1.83 ± 0.36
<i>k</i>	58.15 ± 5.90	-3.39 ± 1.28	48.89 ± 6.16
<i>d</i>	42.39 ± 5.73	102.19 ± 4.65	53.15 ± 7.04
<i>f</i>	1.21 ± 1.07	5.73 ± 2.43	1.28 ± 1.13

ns = not significant; * $P < 0.05$; ** $P < 0.01$.

Original, Monsanto Company) was used. Application of the treatments was performed when the sugarcane plants had five to six true leaves and the eucalyptus shoots averaged 50 cm tall. A 1.5 m wide stationary boom sprayer was used with four XR 110.02 VS spray nozzles spaced every 0.5 m and positioned 0.5 m above the plants. The spray pressure adopted for the equipment was 2.0 kgf cm⁻² with a speed of 3.6 km h⁻¹ and spray volume of 200 L ha⁻¹.

For determination of dry weight, the leaves and stem were separated and then dried in an air-circulating oven at 40 °C. When the material reached a constant weight, it was weighed to 0.001 g precision with a Shimadzu scale (AY220).

2.2 Gas exchange methods

Assessments of gas exchange were carried out at 15 DAA in fully expanded leaves located on the middle third of the eucalyptus plants and in the first completely expanded leaf with a visible auricle in sugarcane.¹³ Analyses for doses above 180 g a.e. ha⁻¹ in eucalyptus were not carried out due to the high level of toxicity caused by this dose of glyphosate. A photosynthesis open system instrument with a CO₂ and water vapor infrared analyzer (IRGA, model LI-6400, Li-Cor) was used. The differences between the CO₂ concentration and water vapor values present in the chamber with and without the samples allowed determination of CO₂ concentration and water vapor released (transpiration) and assimilated (CO₂ assimilation) via the stomata. Gas exchange measures consisted of

the CO₂ assimilation rate (*A*, $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), transpiration rate (*E*, mmol water vapor m⁻² s⁻¹) and stomatal conductance (*g_s*, mol m⁻² s⁻¹). These variables were calculated by the data analysis software of the photosynthesis measuring equipment, which uses the general equation of gas exchanges.¹⁴

2.3 Glyphosate, aminomethylphosphonic acid (AMPA) and shikimate analysis

For the glyphosate, AMPA and shikimic acid analyses, oven-dried plant samples were crushed in a mortar with liquid nitrogen.¹⁵ A 100 mg sample of the crushed matter was placed in a centrifuge tube with 10 mL of acidified water (pH = 2.5). The tubes were then subjected to an ultrasonic bath at an ultrasonic frequency of 42 kHz for 30 min, and centrifuged at 4000 *g* for 10 min at 20 °C. The supernatant was collected and filtered in a Millex HV filter, 0.45 μm , with a 13 mm Durapore membrane and kept in an amber glass vial for further quantification of compounds with an LC-MS/MS system using high-performance liquid chromatography (Shimadzu, model Proeminence UFLC, equipped with two LC-20 AD pumps, an SIL-20 AC auto-injector, a DGU-20A5 degasser, CBM-20A system controller and CTO-20 AC oven) and a hybrid triple quadrupole mass spectrometer (Triple Quad 4500, AB SCIEX).

2.4 Chlorophyll analyses

To determine the chlorophyll content, two leaves were collected, at 15 DAA and 30 DAA for the two species. The chlorophyll content




Figure 3. CO₂ assimilation rate (A) (A, D), stomatal conductance (g_s) (B, E) and transpiration rate (E) (C, F) in sugarcane leaves (A–C) and eucalyptus leaves (D–F) 15 DAA of glyphosate. Data and statistics for this figure are provided in Table 3.

Table 3. Parameters for the regression equations and F values for CO_2 assimilation rate, stomatal conductance and transpiration rate in sugarcane and eucalyptus leaves 15 DAA of glyphosate. Values represent regression parameters $\pm$ standard error. Dose–response curves from these data are plotted in Fig. 3

	Dry weight (% of control)		
	Sugarcane		
	CO_2 assimilation	Stomatal conductance	Transpiration rate
Hypothesis $f \neq 0$	13.95**	6.98*	8.21*
Regression	25.27**	7.35*	6.91**
Model	Model 1	Model 1	Model 1
R²	0.98	0.98	0.97
Constants			
g	-1.82 ± 0.45	-1.35 ± 0.58	-1.06 ± 1.06
b	1.59 ± 0.16	1.71 ± 0.25	1.52 ± 0.28
k	6.54 ± 1.98	0.04 ± 0.01	0.50 ± 0.41
d	3.50 ± 1.70	0.03 ± 0.01	0.73 ± 0.18
f	4.09 ± 2.01	0.04 ± 0.02	0.88 ± 0.97
	Eucalyptus		
	CO_2 assimilation	Stomatal conductance	Transpiration rate
Hypothesis $f \neq 0$	15.28**	10.34**	14.74**
Regression	6.64**	4.11*	4.78*
Model	Model 1	Model 1	Model 1
R²	0.98	0.99	0.99
Constants			
g	-2.79 ± 0.32	-2.67 ± 1.61	-2.63 ± 0.03
b	3.47 ± 1.85	9.38 ± 1.77	9.97 ± 1.69
k	0.34 ± 0.51	0.008 ± 0.008	-0.03 ± 0.11
d	2.01 ± 0.45	0.103 ± 0.011	1.77 ± 0.19
f	0.25 ± 0.14	0.009 ± 0.003	0.16 ± 0.04

* $P < 0.05$; ** $P < 0.01$.

determinations (mg g^{-1}) were based on methods and equations¹⁶: chlorophyll a = $(11.25 \times A_{663} - 2.79 \times A_{647})$; chlorophyll b = $(21.5 \times A_{647} - 5.1 \times A_{663})$; carotenoids = $(1000 \times A_{470} - 1.82 \times \text{chlorophyll a} - 85.02 \times \text{chlorophyll b})/190$; where A is the absorbance at the wavelength indicated.

2.5 Statistical analysis

The data were submitted to analysis of variance by the F test and the means were compared by Tukey's test ($p \leq 0.05$). The statistical models used were those of Brain and Cousins¹⁷ and Streibig,¹⁸ as adapted by Velini *et al.*,¹⁰ with the purpose of describing the dose–response curves with growth stimulation (Model 1) or without growth stimulation (Model 2) after application of glyphosate. The models were fitted by SAS[®] software,¹⁹ and the graphs were plotted by SigmaPlot 12.5 software.

$$\text{Model 1} \quad y = F(x) = \frac{k + fx}{1 + e^{bgx^b}} + d$$

$$\text{Model 2} \quad y = F(x) = \frac{k}{1 + e^{bgx^b}} + d$$

where $y = F(x)$ is the treatment output; x is the herbicide dosage; $k + d$ is the estimated $F(x)$ for the control treatment (without herbicide application); d is the estimated $F(x)$ for herbicide dose causing

maximum inhibition; b determines the way in which $F(x)$ decreases with dose; $g = -\ln(\text{ED}_{50})$, where ED_{50} is defined as the dose that gives 50% of the total achievable effect. The models used are different for only one additional coefficient, which multiplies the independent variable (x), allowing calculation of the increased sum of squares due to regression, including coefficient f with a value different from zero. Therefore, it is possible to test the increase in the model sum of squares resulting from the inclusion of the f constant, with a single degree of freedom. When the increase in the model sum of squares was significant, the hypothesis $f = 0$ was rejected, the occurrence of growth stimulation was accepted, then the complete model (Model 1) was used.¹⁰ When the mean square value was not significant, the hypothesis $f = 0$ was accepted, and it was concluded that there was no growth stimulation at low doses, a standard sigmoid model (Model 2) was used for the data as by Velini *et al.*¹⁰

For the variables of glyphosate and shikimic acid concentrations, data were adjusted by linear regression ($y = a + bx$) as a function of the glyphosate dose, where y is the treatment output, x the herbicide dose, a the intercept when $x = 0$ and b the slope.

3 RESULTS AND DISCUSSION

3.1 Effect on dry matter accumulation

Low doses of glyphosate produced an increase of dry weight of leaves, stems and total shoots in both experiments with both plant species (Figs 1 and 2; Tables 1 and 2). In the first experiment, in which measurements were taken at 40 DAA (Figs 1A and 2A; Tables 1 and 2), glyphosate doses of 7.8 and 7.4 g a.e. ha^{-1} caused increases of 28.8 and 35.3% in total dry weight in sugarcane and eucalyptus, respectively. Whether considering leaf, stem or total shoot dry weight for sugarcane or eucalyptus, the range of glyphosate doses for the maximal hormetic amplitude was a narrow one of only ca 5 to 9 g a.e. ha^{-1} . In the second experiment in which samples were taken at 60 DAA (Figs 1B and 2B; Tables 1 and 2), glyphosate doses of 14.5 and 19 g a.e. ha^{-1} caused increases of 29.4 and 13.1% in total dry weight in sugarcane and eucalyptus, respectively. In the case of sugarcane, the hormetic effect on stem dry weight at 60 DAA (30.3% stimulation) contributed most to the total shoot effect, but this was not the case at 40 DAA. With eucalyptus, the hormetic effects on stem and leaves were similar. For sugarcane, the hormetic effect was higher at 60 DAA than at 40 DAA. But, for eucalyptus, the effect was less at 60 DAA than at 40 DAA.

Previous studies of glyphosate-caused hormesis with eucalyptus and sugarcane have given some similar results with respect to the hormetic dose ranges, but the magnitudes of the effects have been different. In the study of Velini *et al.*¹⁰ with eucalyptus at 60 DAA, the hormetic dose for dry weight, depending on the plant part, ranged from 2 to 4 g a.e. ha^{-1} , whereas the leaf dry weight and stem dry weight were enhanced 60 and 53%, respectively. In that previous study, the root dry weight was enhanced 111% with 1.9 g a.e. ha^{-1} of glyphosate, and the net increase in dry weight of the entire plant was 69.8% with 2.6 g a.e. ha^{-1} of glyphosate. So, there was considerable net increase in dry weight as a result of treatment with a dose of glyphosate equivalent to less than 1% of the recommended dose for weed management. Low glyphosate doses, between 3.6 and 7.2 g a.e. ha^{-1} , increase leaf dry weight of eucalyptus about 20%.²⁰ The dry weight of sugarcane shoots was increased about 35% at 21 DAA by 7.2 to 36 g a.e. ha^{-1} glyphosate,¹³ whereas Silva *et al.*²¹ found glyphosate at 1.8 g a.e. ha^{-1} to stimulate growth of sugarcane up to 85% for the dry mass of shoots. Our results are similar to these earlier findings.




Figure 4. Glyphosate and shikimic acid in sugarcane (A) and eucalyptus (B) leaves 60 DAA of glyphosate. Data and statistics for this figure are provided in Table 4.

Glyphosate hormesis has been observed in both weed and crop species with optimal hormetic doses usually varying from 1.8 to 36 g a.e. ha⁻¹.^{9,10} The optimal dose of glyphosate for hormesis differs with plant species, time after treatment, what is measured, age and physiological status of the plants, and environmental factors.⁹ For example, in a field study with *Brachiaria brizantha*, the optimal dosage for hormetic increases in biomass and plant height at 15 DAA was smaller than the optimal hormetic dose at 30 DAA.²² With coffee plants,²³ there was an 18% increase in stalk diameter, 31% in leaf dry weight and 27% in total dry weight in experiments at 60 DAA with doses of glyphosate near 500 g a.e. ha⁻¹, but glyphosate hormesis was only found in plants treated 45 days after transplanting and not at 10 days after transplanting.

The duration of the hormetic effect after treatment is of great interest, because a long-lasting effect can result in yield increases. In greenhouse studies with barley plants, an increased growth rate in the first week after spraying with glyphosate doses below 60 g a.e. ha⁻¹ was observed.²⁴ A hormetic effect was maintained for 6 weeks after spraying, but was lost after this time and did not result in a yield increase. With sugarcane and eucalyptus, we found that hormesis was maintained for more than 8 weeks, and that, in sugarcane, the effect was greater after 8 weeks than after 5–6 weeks (Table 1).

3.2 Effect of low glyphosate doses on gas exchange

Ultimately, most dry weight increase is due to carbon fixation which can be determined by infrared gas analysis of CO₂ uptake. In the first experiment, the gas exchange evaluation conducted 15 DAA in sugarcane plants had an increase of 99.4% in the CO₂ assimilation rate, compared to the control, for glyphosate applications at 6.1 g a.e. ha⁻¹ (Fig. 3A; Table 3). With this dose, the CO₂ assimilation rate achieved maximum levels of 19.4 µmol m⁻² s⁻¹, almost twice that of the control, which was only 9.6 µmol m⁻² s⁻¹. In eucalyptus plants, the CO₂ assimilation rate increased maximally

Table 4. Parameters for the regression equations and *F* values for glyphosate and shikimic acid in sugarcane and eucalyptus leaves 60 DAA of glyphosate. Values represent regression parameters ± standard error. These data are plotted in Fig. 4

	Concentration (µg g ⁻¹ of dry mass)	
	Sugarcane	
	Glyphosate	Shikimic acid
<i>F</i> values	12.74**	6.06**
<i>R</i>²	0.99	0.894
Constants		
<i>a</i>	0.0010 ± 0.00003	0.0014 ± 0.0004
<i>b</i>	-0.0100 ± 0.0045	1.8187 ± 0.0585
	Eucalyptus	
	Glyphosate	Shikimic acid
	Glyphosate	Shikimic acid
<i>F</i> values	41.81**	3.08*
<i>R</i>²	0.98	0.96
Constants		
<i>a</i>	0.018 ± 0.001	0.062 ± 0.005
<i>b</i>	-0.231 ± 0.102	14.048 ± 0.628
ns = not significant; * <i>P</i> < 0.05; ** <i>P</i> < 0.01; <i>y</i> = <i>ax</i> + <i>y</i> ₀ .		

with glyphosate applications at 11.6 g a.e. ha⁻¹, with an increase of 98.6% compared to the control (Fig. 3D; Table 3). The processes that would cause an increased photosynthetic activity in plants treated with low doses of glyphosate are not clearly understood. The stimulation of photosynthesis in barley plants was reported when exposed to low doses of glyphosate (11 to 45 g a.e. ha⁻¹), and this effect remained until harvest.¹² According to those authors, changes in carbon fixation rate, shikimic acid content and carbohydrate translocation may influence the occurrence of these phenomena.

Table 5. Parameters for the regression equations and *F* values for chlorophyll a and b and total in sugarcane leaves 15 and 30 DAA of glyphosate. Values represent regression parameters $\pm$ standard error. Dose–response curves from these data are plotted in Fig. 5A

	Dry weight (% of control)		
	Sugarcane 15 days after treatment		
	Chlorophyll a	Chlorophyll b	Chlorophyll total
Hypothesis $f \neq 0$	19.04**	90.48**	13.96**
Regression	96.70**	30.24**	33.10**
Model	Model 1	Model 1	Model 1
R^2	0.98	0.99	0.99
Constants			
<i>g</i>	-2.98 ± 0.17	-4.75 ± 0.25	-2.56 ± 0.51
<i>b</i>	3.63 ± 0.88	3.32 ± 1.11	1.72 ± 0.23
<i>k</i>	8.13 ± 1.00	0.93 ± 0.31	7.97 ± 1.44
<i>d</i>	1.84 ± 0.99	2.69 ± 1.15	5.13 ± 2.77
<i>f</i>	0.53 ± 0.27	0.07 ± 0.02	1.21 ± 0.84
	Sugarcane 30 days after treatment		
	Chlorophyll a	Chlorophyll b	Chlorophyll total
	Chlorophyll a	Chlorophyll b	Chlorophyll total
Hypothesis $f \neq 0$	5.24*	7.15*	6.21*
Regression	74.99**	11.18**	22.68**
Model	Model 1	Model 1	Model 1
R^2	0.99	0.99	0.99
Constants			
<i>G</i>	-3.22 ± 0.61	-2.63 ± 0.41	-3.03 ± 0.58
<i>B</i>	1.68 ± 0.24	1.66 ± 0.17	1.65 ± 0.22
<i>K</i>	5.93 ± 0.98	1.42 ± 0.24	7.58 ± 1.31
<i>D</i>	4.39 ± 2.15	1.92 ± 0.99	6.02 ± 2.82
<i>F</i>	0.45 ± 0.37	0.25 ± 0.13	0.73 ± 0.56

* $P < 0.05$; ** $P < 0.01$.

Stomatal conductance assessed 15 DAA in eucalyptus leaves indicated an increase with application of low doses of glyphosate (Fig. 3E; Table 3). For doses of 11.4 g a.e. ha^{-1} , stomatal conductance reached levels of $0.206 \text{ mol m}^{-2} \text{ s}^{-1}$, corresponding to an 82.7% increase in relation to the control. Similarly, stomatal conductance in sugarcane increased with a dose of 3.4 g a.e. ha^{-1} (Fig. 3B; Table 3). The maximum stomatal conductance value observed was $0.123 \text{ mol m}^{-2} \text{ s}^{-1}$, this value corresponding to an increase of 82.7% with respect to the control.

The sugarcane transpiration rate increased maximally (80.6%) with glyphosate application at 3.4 g a.e. ha^{-1} compared to the control, reaching $2.3 \text{ mmol m}^{-2} \text{ s}^{-1}$ (Fig. 3C; Table 3). The transpiration rate of eucalyptus increased with glyphosate application (Fig. 3F; Table 3) with a maximum value of $3.3 \text{ mmol m}^{-2} \text{ s}^{-1}$ (86.1% higher than the control) found at a glyphosate application rate of 11.2 g a.e. ha^{-1} .

The doses that caused increases in gas exchange variables were all below 10 g a.e. ha^{-1} . In glyphosate-treated eucalyptus plants, CO_2 uptake, stomatal conductance and transpiration rate diminished when doses above 43.2 g a.e. ha^{-1} were applied.²⁵

Stomatal movement is the main mechanism of control of gas exchange in all but primitive plants, because virtually all CO_2 influx and water efflux occur through the stomata. Control of gas exchange is a complex process because plants face the dilemma associated with a proper balance between CO_2 uptake and water loss by maintaining a stomatal aperture that avoids water stress

while maximizing carbon fixation, a balance that is problematic under even mild water stress. Photosynthesis depends on flow of CO_2 into the cell, and this CO_2 flow depends on the stomatal opening.²⁶ Rubisco, the enzyme responsible for assimilation of CO_2 , is a highly inefficient enzyme because O_2 competes with CO_2 , resulting in photorespiration, a process that wastes ATP and NADPH from the light reactions of photosynthesis. Thus, the higher the CO_2 to O_2 ratio, the more efficiently Rubisco can function. Thus, increased stomatal conductance has a beneficial relation in the Rubisco activity and, consequently, in photosynthesis.²⁷ High rates of CO_2 uptake have a direct relationship with loss of water by transpiration. Thus, with adequate water, high stomatal conductance leads to high water consumption and a positive growth increase.²⁸

Glyphosate was not detected in eucalyptus leaves with doses of 1.8 to 7.2 g a.e. ha^{-1} (Fig. 4B; Table 4) and doses of 1.8 to 36 g a.e. ha^{-1} for sugarcane (Fig. 4A; Table 4). The minimum detection of glyphosate in spiked untreated leaf tissue was $0.25 \mu\text{g g}^{-1}$ dry weight, with a recovery of about 85% for both species. The glyphosate levels found at the three highest rates of glyphosate application were about ten-fold higher than those in sugarcane. AMPA was found only in eucalyptus and only at very low concentrations at the three highest glyphosate treatments. Considering that the herbicide was only quantified in the leaves, it is possible that a significant part of the herbicide had been translocated to the roots or stem by 60 DAA. With doses of 18 g a.e. ha^{-1} and higher, the glyphosate levels in the leaves increased progressively up to the dose of 360 g a.e. ha^{-1} , at which the glyphosate concentration for sugarcane was $0.35 \mu\text{g g}^{-1}$ dry weight and that for eucalyptus was $6.3 \mu\text{g g}^{-1}$ dry weight (Fig. 4).

At 60 DAA of 360 g a.e. ha^{-1} glyphosate, the shikimic acid concentration in the sugarcane leaves increased 23.6% compared to the control (Fig. 4A), reaching a level of about $2.3 \mu\text{g g}^{-1}$ dry weight. In eucalyptus leaves, for the same dosage, the shikimic acid levels increased 2.5 times in relation to the control, reaching values of about $36.1 \mu\text{g g}^{-1}$ dry weight. Shikimic acid accumulation is usually directly associated with glyphosate concentrations.²⁹ The dose–response curves of the glyphosate and shikimate contents were similar in both species, although the control levels of shikimate were higher in eucalyptus than in sugarcane.

The finding that no glyphosate or glyphosate-induced shikimate levels were found at the doses of glyphosate that stimulated growth is surprising. However, glyphosate and shikimate concentrations were determined at 60 DAA, and the carbon assimilation was measured at 15 DAA. Measureable glyphosate and an increase in shikimate could have been detected at 15 DAA and lost later due to translocation and/or degradation. Glyphosate-elevated shikimate levels are known to be reduced or lost over time.³⁰

An increase in the content of chlorophylls and carotenoids occurred in sugarcane plants that received low doses of glyphosate (Table 5; Fig. 5). However, there was no effect of herbicide glyphosate doses on chlorophyll in eucalyptus (data not shown). Many studies have reported reductions in chlorophyll content in plants treated with phytotoxic doses of glyphosate.^{31–33} However, subtoxic dose effects of glyphosate on chlorophyll content in plants are not found in the literature.

4 CONCLUSIONS

Application of low doses of glyphosate, in the range of 5.8 to 19 g a.e. ha^{-1} , increased leaf, stem and total dry weight in sugarcane and eucalyptus plants at 40 and 60 DAA. At 15 DAA, CO_2




Figure 5. Chlorophyll a and b and total chlorophyll contents in sugarcane leaves 15 (A) and 30 (B) DAA of glyphosate. Data and statistics for this figure are provided in Table 5.

assimilation, stomatal conduction and transpiration rates were stimulated by sublethal doses, ranging from 3.4 to 11.6 g a.e. ha^{-1} . The hormetic effect on carbon assimilation is likely to be responsible for the later hormetic effect on dry weight increases. The dose–response curves for glyphosate shikimate content were similar for sugarcane and eucalyptus, with both chemicals only increasing at doses above about 100 g a.e. ha^{-1} .

ACKNOWLEDGEMENTS

We are grateful for the financial support from São Paulo Research Foundation (FAPESP) and Brazilian National Council for Scientific and Technological Development (CNPq).

REFERENCES

- Duke SO, Cedergreen N, Velini ED and Belz RG, Hormesis: is it an important factor in herbicide use and allelopathy? *Outlooks Pest Manag* **17**:29–33 (2006).
- Duke SO and Powles SB, Glyphosate: a once-in-a-century herbicide. *Pest Manag Sci* **64**:319–325 (2008).
- Tan S, Evans R and Singh B. Herbicidal inhibitors of amino acid biosynthesis and herbicide-tolerant crops. *Amino Acids* **30**:195–204 (2006).
- Rippert P, Scimemi C, Dubald M and Matringe M, Engineering plant shikimate pathway for production of tocotrienol and improving herbicide resistance. *Plant Physiol* **134**:92–100 (2004).
- Herrmann KM and Weaver LM, The shikimate pathway. *Annu Rev Plant Biol* **50**:473–503 (1999).
- Dewick PM, The biosynthesis of shikimate metabolites. *Nat Prod Rep* **15**:17–58 (1998).
- Herrmann KM, The shikimate pathway: early steps in the biosynthesis of aromatic compounds. *Plant Cell* **7**:907–919 (1995).
- Calabrese EJ, Bachmann KA, Bailer AJ, Bolger M, Borak J, Cai L et al., Biological stress response terminology: integrating the concepts of adaptive response and preconditioning stress within a hormetic dose–response framework. *Toxicol Appl Pharmacol* **222**:122–128 (2007).
- Belz RG and Duke SO, Herbicides and plant hormesis. *Pest Manag Sci* **70**:698–707 (2014).
- Velini ED, Alves E, Godoy MC, Mechede DK, Souza RT and Duke SO, Glyphosate at low doses can stimulate plant growth. *Pest Manag Sci* **64**:489–496 (2008).
- Cedergreen N and Olesen CF, Can glyphosate stimulate photosynthesis? *Pestic Biochem Physiol* **96**:140–148 (2010).
- Carbonari CA, Gomes GLGC, Velini ED, Machado RF, Simões PS and Macedo GC, Glyphosate effects on sugarcane metabolism and growth. *Am J Plant Sci* **5**:3585–3593 (2014).
- Martins MT, de Souza WR, da Cunha BA, Basso MF, de Oliveira NG, Vinecky F et al., Characterization of sugarcane (*Saccharum* spp.) leaf senescence: implications for biofuel production. *Biotechnol Biofuels* **9**:153 (2016).
- Von Caemmerer S and Farquhar GD, Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta* **153**:376–387 (1981).
- Gomes GLGC, Carbonari CA, Velini ED, Trindade MLB and Silva JRM, Extraction and simultaneous determination of glyphosate, AMPA and compounds of the shikimic acid pathway in plants. *Planta Daninha* **33**:295–304 (2015).
- Lichtenthaler HK, Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol* **148**:350–385 (1987).
- Brain P and Cousens R, An equation to describe dose–responses where there is stimulation of growth at low doses. *Weed Res* **29**:93–96 (1989).
- Streibig JC, Models for curve-fitting herbicide dose response data. *Acta Agric Scand* **30**:59–64 (1980).
- SAS Institute, *SAS User's Guide: Statistics, Version 9.2*, 6th edition. SAS Institute, New York (2008).
- Pereira FCM, Nepomuceno MP, Pires RN, Pareira MC and Alves PLCA, Response of eucalyptus (*Eucalyptus urograndis*) plants at different doses of glyphosate. *J Agric Sci* **5**:66–74 (2013).
- Silva MA, Aragão, Barbosa MA, Jeronimo, EM and Carlin SD, Efeito hormético de glyphosate no desenvolvimento inicial de cana-de-açúcar. *Bragantia* **68**:973–978 (2009).
- Nascentes RF, Fagan EB, Soares LH, Oliveira LH, Oliveira CB and Brunelli MC, Hormesis de glyphosate em *Brachiaria brizantha* cv. Marandu. *Cerrado Agrobiociências* **6**:55–64 (2015).
- Carvalho LB, Alves PLCA and Duke SO, Hormesis with glyphosate depends on coffee growth stage. *Ann Brazilian Acad Sci* **85**:813–821 (2013).
- Cedergreen N, Is the growth stimulation by low doses of glyphosate sustained over time? *Environ Pollut* **156**:1099–1104 (2008).
- Machado AFL, Ferreira LR, Santos LDT, Ferreira FA, Viana RG, Machado MS et al., Eficiência fotossintética e uso da água em plantas de

- eucalipto pulverizadas com glyphosate. *Planta Daninha* **28**:319–327 (2010).
- 26 Messinger SM, Buckley TN and Mott KA, Evidence for involvement of photosynthetic processes in the stomatal response to CO₂. *Plant Physiol* **140**:771–778 (2006).
- 27 Long SP, Zhu X-G, Naidu SL and Ort DR, Can improvement in photosynthesis increase crop yields? *Plant Cell Environ* **29**:315–330 (2006).
- 28 Farquhar GD and Thomas DS, Stomatal conductance and photosynthesis. *Annu Rev Plant Physiol* **33**:317–345 (1982).
- 29 Buehring NW, Massey JH and Reynolds DB, Shikimic acid accumulation in field-grown corn (*Zea mays*) following simulated glyphosate drift. *J Agric Food Chem* **55**:819–824 (2007).
- 30 Lydon J and Duke SO, Glyphosate induction of elevated levels of hydroxybenzoic acids in higher plants. *J Agric Food Chem* **36**:813–818 (1988).
- 31 Reddy KN, Rimando AM and Duke SO, Aminomethylphosphonic acid, a metabolite of glyphosate, causes injury in glyphosate-treated, glyphosate-resistant soybean. *J Agric Food Chem* **52**:5139–5143 (2004).
- 32 Kitchen LM, Witt WW and Rieck CE, Inhibition of chlorophyll accumulation by glyphosate. *Weed Sci* **29**:513–516 (1981).
- 33 Lee TT, Effects of glyphosate on synthesis and degradation of chlorophyll in soybean and tobacco cells. *Weed Res* **21**:161–164 (1981).